



UNIVERSITY OF JOHANNESBURG

FACULTY OF SCIENCE

DEPARTMENT OF APPLIED MATHEMATICS

MODULE **APM1B10**
 INTRODUCTION TO DYNAMICS
CAMPUS **APK**

NOVEMBER EXAMINATION

DATE: 05/11/2014

SESSION: 12:30 - 15:30

ASSESSOR

DR JSC PRENTICE

INTERNAL MODERATOR

PROF C M VILLET

DURATION: **$2\frac{1}{2}$ HOURS**

MARKS: **50**

NUMBER OF PAGES: 4 PAGES

INSTRUCTIONS:

CALCULATORS MAY BE USED

ANSWER ALL THE QUESTIONS

SYMBOLS HAVE THEIR USUAL MEANING

PHYSICAL QUANTITIES ARE IN SI UNITS AND ANGLES ARE IN RADIANS

QUESTION 1

(a) Show that

$$\hat{r} \cdot \hat{n} = \frac{v_{\theta} a_n}{v_{\theta} a_r - v_r a_{\theta}}.$$

(b) The equation of the trajectory of a particle is given by

$$\mathbf{r}(t) = t^3 \hat{x} + \sin(2t) \hat{y} \equiv (t^3, \sin(2t)).$$

Find, for $t = \frac{1}{2}$,

- (i) \mathbf{v} and \mathbf{a} ,
- (ii) $\hat{\tau}$ and \hat{n} ,
- (iii) the tangential and normal components of \mathbf{a} ,
- (iv) and the radius of curvature ρ .

[10]

QUESTION 2

Calculate $\int_{\Gamma} \mathbf{a} \cdot d\mathbf{r}$ with $\mathbf{a} = (2xy, x^2, 2z)$, and where the path Γ is defined by

$$\begin{aligned} y &= 2x + 1 \\ z &= x + y^2. \end{aligned}$$

In the integral, the lower limit is $(0, 1, 1)$, and the upper limit is $(1, 3, 10)$. You should find that the value of the integral is 102.

[10]

QUESTION 3

A force $\mathbf{F} = (F_x, F_y)$ acts on a particle, of mass 0.2kg, for 2 seconds. Here, F_x and F_y are constants, not necessarily equal. Assume that \mathbf{F} makes an angle θ with the horizontal axis. Assume that the *horizontal* component of the initial velocity \mathbf{v}_i is half the vertical component. Assume that the final velocity \mathbf{v}_f has the same magnitude as \mathbf{v}_i , but is opposite in direction.

- (a) Determine θ .
- (b) If the horizontal component of \mathbf{v}_i is 20ms^{-1} in the *negative* x -direction, determine F_x and F_y .

[10]

QUESTION 4

Calculate the directional derivative $\frac{\partial \psi}{\partial s}$ of

$$\psi = yz - x^2$$

in the direction

$$\hat{x} + 2\hat{y} - 3\hat{z}$$

at the point

$$(-2, 1, 0).$$

(a) Use the parameterization $\mathbf{r} = \mathbf{r}_0 + s\hat{s}$.

(b) Use $\nabla\psi$.

[10]

QUESTION 5

A bead of mass m slides along a smooth circular ring from A to B (Figure 1). Calculate the speed at B if the initial speed was v_0 . The angle θ is defined in the figure. Gravity acts vertically downwards.




Figure 1

[10]

INFORMATION

$$\begin{aligned}
 \mathbf{r} &= (x, y) = (r \cos \theta, r \sin \theta) \\
 \mathbf{v} &= v_r \hat{r} + v_\theta \hat{\theta} = v_\tau \hat{\tau} + v_n \hat{n} \\
 &= \dot{r} \hat{r} + r \dot{\theta} \hat{\theta} = v \hat{\tau} \\
 \mathbf{a} &= a_r \hat{r} + a_\theta \hat{\theta} = a_\tau \hat{\tau} + a_n \hat{n} \\
 &= \left(\ddot{r} - r \dot{\theta}^2 \right) \hat{r} + \left(r \ddot{\theta} + 2 \dot{r} \dot{\theta} \right) \hat{\theta} = \dot{v} \hat{\tau} + \left(\frac{v^2}{\rho} \right) \hat{n} \\
 \hat{r} &= (\cos \theta, \sin \theta) & \hat{\theta} &= (-\sin \theta, \cos \theta) \\
 \hat{\tau} &= (\cos \psi, \sin \psi) & \hat{n} &= (-\sin \psi, \cos \psi)
 \end{aligned}$$

$$\mathbf{I} = \int_{t_i}^{t_f} \mathbf{F} dt = m \mathbf{v}_f - m \mathbf{v}_i$$

$$\begin{aligned}
 \frac{\partial \Psi}{\partial s} &= \nabla \Psi \cdot \hat{s} \\
 d\mathbf{r} &= (dx, dy) \text{ in two dimensions} \\
 d\mathbf{r} &= (dx, dy, dz) \text{ in three dimensions}
 \end{aligned}$$

$$\begin{aligned}
 T &= \frac{m(\mathbf{v} \cdot \mathbf{v})}{2} = \frac{mv^2}{2} \\
 W &= \int_{\Gamma} \mathbf{F} \cdot d\mathbf{r} \\
 \mathbf{F} &= -\frac{GMm}{r^2} \hat{r}
 \end{aligned}$$